

9700196

THE UNITED SHAYES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Pioneer Hi-Bred International, Inc.

ILCCOR, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE AIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR ORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT OF BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PHDG1'

In Destinour Mercus, I have hereunto set my hand and caused the seal of the Hant Anticip Arotection Office to be affixed at the City of Washington, D.C. this fifth day of February, in the year of our Lord two thousand one.

alanktost

Acting Commissioner Plant Variety Protection Office Agricultural Marketing Service Secretary of Agriculture

SD-470 (04-95) (Previous editions are to be destroyed)

14A. Exhibit A. Origin and Breeding History

Pedigree: **#G**50/PHP02)X3K132311

JUS 125/00

Pioneer Line PHDG1, Zea mays L., a yellow corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross EEG50 X PHP02 using the pedigree method of breeding. The progenitors of PHDG1 are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing and selection were practiced within the above F1 cross for 7 generations in the development of PHDG1 at Eau Claire, Wisconsin. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Eau Claire, Wisconsin, as well as other Pioneer research stations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations made for uniformity.

PHDG1 has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line PHDG1 has been increased both by hand and in isolated fields with continued observations for uniformity and stability throughout development, and for 3 generations

during the final stages of inbred development and seed multiplication.

No variant traits have been observed or are expected in PHDG1.

The criteria used in the selection of PHDG1 were yield, both per se and in hybrid combinations; kernel size, especially important in production; ability to germinate in adverse conditions; number of tillers, especially important in production because having numerous tillers increases hybrid production costs spent on detasseling; disease and insect resistance; pollen yield; tassel size and pollen shed duration.

SMS 6/27/00

Sugar Francis Land

DEVELOPMENTAL HISTORY FOR PHDG1

Season/Ye	ear	Inbreeding	Level
Summer 19	985	FO	
Winter 19	985	F1	
Summer 19	986	F2*	
Winter 19	986	F3*	
Summer 19	987	F4*	
Summer 19	988	F5*	
Summer 19	989	F6*	
Summer 19	990	F7*	
Summer 19	991	F8*	
Winter 19	991	F9*	Ļ

^{*}PHDG1 was selfed and selected through F8 generation. **PHDG1 was selfed and ear-rowed for F9 generation.

Exhibit B Novelty Statement

Variety PHDG1 mostly resembles the Pioneer Hi-Bred International, Inc. proprietary inbred line PHR31 (PVP Certificate No.9200090).

Variety PHDG1 reaches 50% pollen shed (GDUSHD) earlier (1325 GDU's vs 1369 GDU's) than PHR31 (Exhibit D).

Variety PHDG1 reaches 50% silking (GDUSLK) earlier (1368 GDU's vs 1399 GDU's) than PHR31 (Exhibit D).

Variety PHDG1 has a longer tassel peduncle length (19.3 cm vs 14.8 cm) than PHR31 (Table 1A, 1B).

Variety PHDG1 has a wider tassel branch angle (41.7 degrees vs 24.0 degrees) than PHR31 (Table 1A, 1B).

Variety PHDG1 has a thicker kernel thickness (5.1 mm vs 4.3 mm) than PHR31 (Table 1A, 1B).

Exhibit B Novelty Statement Tables

Table 1A Data from Johnston, IA in 1998 are supporting evidence for differences between PHDG1 and PHR31. Locations had different environmental conditions

2001				ö									
Statio	Station loc Year	Year	Traits	Variety-	Variety-	Count-	Count- Mean-		Mean-	Wean	DF	L-Value	Prob (2-
				~	2	-	N.		N	斯	Pooled	Pooled	tail) Pooled
PP	20N	1998	kernel thickness (mm)	PHDG1		5	5	4.8	4.2	9.0	8	2.12	0.067
ᆽ	92	1998	kernel thickness (mm)	PHDG1	PHR31	2	5	5.4	4.4	1.0	œ	2.89	0.020
ΑD	20N	1998	tassel branch angle (degrees)	PHDG1		ည	5	35.4	26.0	9.4	æ	3.04	0.016
ᆽ	95	1998	tassel branch angle (degrees)	PHDG1		2	ည	48.0	22.0	26.0	œ	5.15	0.001
AD	20N	1998	tassel peduncle length (cm)	PHDG1		2	2	18.6	14.2	4.4	ω	4.02	0.004
亐	95	1998	tassel peduncle length (cm)	PHDG1		5	5	20.0	15.4	4.6	œ	9.05	0.000
				The state of the s	Commence of the Commence of th	THE PERSON NAMED IN COLUMN 2 IN COLUMN 2	The second secon	THE PERSON NAMED IN COLUMN 1	The second second second	THE PERSON NAMED IN COLUMN 1	THE RESIDENCE OF THE PERSON.	The second secon	ARRAY CANADA TANADA TAN

Table 1B Summary data from Johnston, IA across 1998 are supporting evidence for differences between PHDG1 and PHR31 locations had different environmental conditions.

Profit (2- tail) Pooled	0.003	0.000	0.000
t-Value Prob (2- Pooled tail) Pooled	3.39	4.98	6.95
Pooled	18	17.7 18	18
02864	9.0	17.7	4.5
(ean-2	4.3	24.0	14.8
nte Count-Mean+1 Mean-2 Mean 2 Diff	5.1	41.7	19.3
Count- 2	10	10	10
-	10	10	10
Variety-1 Variety ©o	PHR31	PHR31	PHR31
Variety-1	PHDG1	PHDG1	PHDG1
Vear	1998 kernel thickness (mm)	1998 tassel branch angle (degrees)	1998 tassel peduncle length (cm)

Exhibit C (Corn: Maize)

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

Objective Description of Variety Corn (Zea mays L.)

Name of Applicant (s)		Variety Seed Source	Variety Nam	Variety Name or Temporary Designation				
Pioneer 1	r Hi-Bred International, Inc.				PHDG1			
Address (Str	reet & No., o	or RFD No., City, State, Zip Coo	de and Country	FOR OFFICIAL USE	3			
7301 NW	/ 62 nd Av	enue, P.O. Box 85,						
Johnston	ı, Iowa 5	0131-0085		PVP0 Number				
Place the ap	propriate nui	mber that describes the varietal	characters typical of this inbred	variety in the spaces be	low. Right justify whole			
numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits								
designated by an '*' are considered necessary for an adequate variety description and must be completed.								
COLOR CHOICES (Use in conjunction with Munsell color code to describe all color choices: describe #25 and #26 in Comments section):								
01=Light Green 06=Pale Yellow 11=Pink		11=Pink	16=Pale Purple	21=Buff				
02=Medium	Green	07=Yellow	12=Light Red	17=Purple	22=Tan			
03=Dark Gre	een	08=Yellow Orange	13=Cherry Red	18=Colorless	23=Brown			
04=Very Da	rk Green	09=Salmon	14=Red	19=White	24=Bronze			
05=Green-Y	ellow	10=Pink-Orange	15=Red & White	20=White Capped	25=Variegated (Describe)			
					26=Other (Describe)			
STANDARI								
•	,	background and maturity) of the	-	•	The state of the s			
Yellow Dent	t Families:		Yellow Dent (Unrelated	l): Sweet (Corn:			
Family Members Co10		Co109, ND246,	C13, I	owa5125, P39, 2132				
B14	14 CM105, A632, B64, B68 Oh7, T232,							
B37			n:					
B73	N192, A679	, B73, NC268	W18BN	SG153	33, 4722, HP301, HP7211			
C103	Mo17, Va10	2, Va35, A682						
Oh43	A619, MS71	I, H99, Va26	White Dent:	Pipecor	n:			
WF9	W64A, A55	4, A654, Pa91	C166, H105, Ky228	Mo15	W, Mo16W, Mo24W			

Ceres/worddata/doug/96pvp

EXHIBIT C: PHDG1

Application Variety Data

1. TYPE: (describe intermediate types in Comments section): Standard Variety Name: W64A 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental 2. REGION WHERE DEVELOPED IN THE U.S.A.: Standard Seed Source: AMES 19291 1=Northwest 2=Northcentral 3=Northeast 4=Southeast 5=Southcentral 6=Southwest 7=Other 3. MATURITY (In Region of Best Adaptability; show Heat Unit formula in 'Comments' section) DAYS **HEAT UNITS** DAYS **HEAT UNITS** 068 1,341.8 From emergence to 50% of plants in silk 066 1,317.4 1,289.0 066 From emergence to 50% of plants in pollen 065 1,291.8 004 0.096.8 From 10% to 90% pollen shed 005 0,113.6 From 50% silk to optimum edible quality 068 1,274.2 From 50% silk to harvest at 25% moisture 066 1,268.6 4. PLANT: Standard Sample Standard Sample Deviation Size Deviation Size 208.6 cm Plant Height (to tassel tip) 10.14 05 165.4 05.37 05 078.2 cm Ear Height (to base of top ear node) 10.92 05 060.2 03.90 05 015.6 cm Length of Top Ear Internode 01.14 05 012.1 01.14 05 0.0 Average Number of Tillers 00.02 05 0.0 00.01 05 1.2 Average Number of Ears per Stalk 00.45 05 05 1.0 00.00 2 Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark 5. LEAF: Standard Sample Standard Sample Size Deviation Deviation Size 08.2 cm Width of Ear Node Leaf 00.38 05 08.6 00.73 05 81.4 cm Length of Ear Node Leaf 04.33 05 62.5 06.48 05 06 Number of leaves above top ear 00.26 05 00.68 05 05 39 Degrees Leaf Angle (measure from 2nd leaf above ear 05.28 05 04.28 05 44 at anthesis to stalk above leaf) 03 Leaf Color (Munsell code) 5GY 4/4 03 5GY 4/4 1 Leaf Sheath Pubescence (Rate on scale from 1=none to 9=like peach fuzz) 1 7 Marginal Waves (Rate on scale from 1=none to 9=many) 6 6 Longitudinal Creases (Rate on scale from 1=none to 9=many) 6 6. TASSEL: Standard Sample Standard Sample Deviation Size Deviation Size 14 **Number of Primary Lateral Branches** 02.21 05 05 05 01.58 59 Branch Angle from Central Spike 05 22.15 05 24 08.19 56.8 cm Tassel Length (from top leaf collar to tassel tip) 04 01.57 04 50.1 03.76 7 Pollen Shed (rate on scale from 0=male sterile to 9=heavy shed) 7 07 Anther Color (Munsell code) 10Y 8.5/8 07 10Y 8.5/8 01 Glume Color (Munsell code) 5GY 6/6 01 5GY 6/6 1 Bar Glumes (Glume Bands): 1=Absent 2=Present

Page 1

Standard Variety Data

JUS 1/27/00

Application	Nariety Data PHD)G1	Page 2			Standa	rd Inbred	Data
7a. EAR	(Unhusked Data):				ORP 6/6			
11 -01	Silk Color (3 days after emerge	ence) (Mur	isell code)	I.	7.5Y66	<u>07</u>	2.5G	<u> 196</u>
02	Fresh Husk Color (25 days aft	er 50% silki	ing) (Munsell code)	1	5GY56	01	5GY	
21	Dry Husk Color (65 days after	50% silking) (Munsell code)		2.5Y84	21	2.5Y8	
1	Position of Ear at Dry Husk St	age: 1= Upr	right 2= Horizontal	3≕ Pendant		<u>3</u>		
5	Husk Tightness (Rate of Scale	from 1=ve	ry loose to 9=very	tight)		<u>6</u>		
2	Husk Extension (at harvest): 1	=Short (ear	s exposed) 2=Med	ium (<8 cm)		<u>2</u>		
	3=Long (8-10 cm beyond ear t	ip) 4=Very I	Long (>10 cm)					
7b. EAF	(Husked Ear Data):			Standard	Sample	Star	ndard	Sampl
	,			Deviation	Size	Dev	iation	Size
<u>15.1</u>	cm Ear Length			01.02	<u>05</u>	<u>12.7</u> <u>C</u>	00.71	<u>05</u>
<u>43.0</u>	mm Ear Diameter at mid-point			02.36	<u>05</u>	41.8 C	1.38	<u>05</u>
117.2	gm Ear Weight			14.00	<u>05</u>	86.6 C	<u> </u>	<u>05</u>
<u>15</u>	Number of Kernel Rows			00.56	<u>05</u>	<u>17.1</u> C	1.60	<u>05</u>
2	Kernel Rows: 1=Indistinct 2=D	istinct				<u>2</u>		
<u>1</u>	Row Alignment: 1=Straight 2=	Slightly Cur	ved 3=Spiral			1		
<u>14.8</u>	cm Shank Length			02.95	<u>05</u>	<u>09.5</u> <u>0</u>	00.72	<u>05</u>
<u>1</u>	Ear Taper: 1=Slight 2= Averag	e 3=Extren	пе			<u>2</u>		
8. KERN	EL (Dried)			Standard	Sample	Standa	ard	Sampl
				Deviation	Size	Deviati	ion	Size
<u>10.6</u>	mm Kernel Length			00.50	<u>05</u>	<u>09.2</u> <u>0</u>	0.19	<u>04</u>
08.2	mm Kernel Width			00.22	<u>05</u>	<u>07.0</u> 0	0.71	<u>04</u>
<u>05.1</u>	mm Kernel Thickness			00.50	<u>05</u>	04.4 0	<u>0.46</u>	<u>04</u>
<u>45.2</u>	% Round Kernels (Shape Grad	e)		09.50	<u>04</u>	<u>06.0</u> 0	5.98	<u>04</u>
<u>1</u>	Aleurone Color Pattern: 1-Hom	nozygous 2:	=Segregating			1		
<u>07</u>	Aluerone Color (Munsell code)			<u>10</u> °	YR814	<u>07</u>	2.5Y8	<u>312</u>
<u>07</u>	Hard Endosperm Color (Munse	ell code)		<u>10`</u>	YR714	<u>07</u>	<u>10YR</u>	<u>814</u>
<u>03</u>	Endosperm Type:					<u>3</u>		
	1=Sweet (Su1) 2=Extra Sw 4=High Amylose Starch 5=\ 7=High Lysine 8=Super Sw 10=Other	Naxy Starc	h 6=High Protein					
<u>30.4</u>	gm Weight per 100 Kernels (un	sized samp	ile)	03.65	<u>05</u>	<u>19.00</u> <u>0</u>	4.24	<u>04</u>
9. COB:			, ,,,,, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Standard	Sample	St	andard	Samp
				Deviation	Size	De	eviation	Size
<u>26.4</u>	mm Cob Diameter at mid-point			01.82	<u>05</u>	<u>27.4</u>	01.08	<u>04</u>
14	Cob Color (Munsell code)		10R46			<u>14</u>	2.5Y	DEG

Application Variety Data

Page 3

Standard Inbred Data

40 0105405 0		
	ESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant); le ; leave Race or Strain Options blank if polygenic):	ave blank
A. Leaf B	lights, Wilts, and Local Infection Diseases	
	Anthracnose Leaf Blight (Colletotrichum graminicola)	
	Common Rust (Puccinia sorghi)	
	Common Smut (Ustilago maydis)	
	Eyespot (Kabatiella zeae)	
<u>7</u>	Goss's Wilt (Clavibacter michiganense spp. nebraskense)	<u>7</u>
<u>5</u>	Gray Leaf Spot (Cercospora zeae-maydis)	<u>2</u>
	Helminthosporium Leaf Spot (Bipolaris zeicola) Race ———	
<u>5</u>	Northern Leaf Blight (Exserohilum turcicum) Race ———	<u>5</u>
	Southern Leaf Blight (Bipolaris maydis) Race ——	
	Southern Rust (Puccinia polysora)	
<u>4</u>	Stewart's Wilt (Erwinia stewartii)	1
	Other (Specify)	
B. Systen	nic Diseases	
<u>3</u>	Corn Lethal Necrosis (MCMV and MDMV)	2
<u>-</u> <u>5</u>	Head Smut (Sphacelotheca reiliana)	<u>2</u> 9
_	Maize Chlorotic Dwarf Virus (MDV)	_
	Maize Chlorotic Mottle Virus (MCMV)	
	Maize Dwarf Mosaic Virus (MDMV)	
	Sorghum Downy Mildew of Corn (Peronosclerospora sorghi)	
	Other (Specify) ———	
C. Stalk F	Rots	
•	Anthracnose Stalk Rot (Colletotrichum graminicola)	
	Diplodia Stalk Rot (Stenocarpella maydis)	
	Fusarium Stalk Rot (Fusarium moniliforme)	
	Gibberella Stalk Rot (Gibberella zeae)	
	Other (Specify) ——	
D. Ear an	d Kernel Rots	
	Aspergillus Ear and Kernel Rot (Aspergillus flavus)	
	Diplodia Ear Rot (Stenocarpella maydis)	
	Fusarium Ear and Kernel Rot (Fusarium moniliforme)	
<u>4</u>	Gibberella Ear Rot (Gibberella zeae)	<u>4</u>
_	Other (Specify)	-

CLARIFICATION OF DATA IN EXHIBITS C AND D

Please note the data presented in Exhibit C, "Objective Description of Variety," is data collected primarily at Johnston, Iowa plus description information from the maintaining station. The data in Exhibit D, "Additional Description of Variety," is data from comparisons of inbreds grown in the same tests in the adapted growing area of PHDG1.

JH5 00

There are environmental factors that differ from year to year. In 1995, May was wet and August was warmer. In 1996, May was very wet and August was cool with very little heat or drought stress compared to most years. Environmental temperature and precipitation differences during the vegetative and grain fill periods can impact plant and grain traits and be a source of variability. Please see table 3, which summarizes rainfall and growing season temperatures from 1994-1997. The environmental conditions described above could result in larger standard deviations. Also, in particular the tassel branch angle trait is very subject to variability based on how a person measures it. We try to monitor our measuring procedures carefully but measuring the angle from different primary branches could result in higher variability for this trait. Since I was not involved in the data collection of this trait at that time I am not sure what caused the variability for this trait.

Table 3. Average temperatures (Fahrenheit) and rainfall (inches) for central Iowa.

TEMPERATURE

YEAR	MAY	JUN	JULY	AUG	AVERAGE
1994	59.8	70.7	71.9	69.0	67 <i>.</i> 9
1995	56.2	69.4	74.3	76.9	69.2
1996	56.2	69.3	71.3	70.5	66.8
1997	53.5	70.6	74.1	69.6	67.0
AVG	56.4	70.0	72.9	71.5	67.7

RAINFALL

YEAR	MAY	JUN	JULY	AUG	Total
1994	3.67	5.75	1.71	4.18	15.31
1995	5.04	4.19	2.94	2.87	15.04
1996	8.47	4.35	2.51	2.14	17.47
1997	4.32	3.27	4.10	1.36	13.05
AVG	5.38	4.39	2.82	2.64	15.22

ADDITIONAL DESCRIPTION OF PHDG1 INBRED PER SE YIELD TEST COMPARISON OF PHDG1 AND PHR31 EVALUATED OVER YEARS EXHIBIT D.

VARIETY #1 = PHDG1 VARIETY #2 = PHR31

	i tri tri to	ı	0 0 H 0 0	44 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	002206	
SIG	DRP EAR ABS		100. 100. 1 0.	98. 100. 1.	100. 100. 0.	99. 100. 1 0.
#	BAR PLT ABS	97.8 97.1 6 6 7.0	96.7 99.3 13 15 2.6 151.	100.0 100.0 3 3 0.0 9.0	100.0 98.1 2 2 1.9	97.6 98.9 34 36 1.3
IG	GRN APP ABS		5.5 5.6 6.1 861	8.3 7.0 3 1.3 1.3	7.0 7.8 4 4 0.8	6.6 12 19 0.0
រេ % ល	BRT STK ABS	93.1 96.1 7 3.0 3.0	80.0 00.0 1 20.0		87.0 92.3 1 5.3	93.2 97.1 12 12 3.8 3.8
SIG +	STK LDG ABS	90.0 90.5 9.5 9.5	99.4 99.2 1 6 11 0.2	95.1 82.2 4 12.9 216	96.6 85.4 3 111.2	97.6 90.8 14 23 6.9
10% S	STA	5.2 1 3.7 6 1.5	5.7 5.3 3 0.4 808	6.8 5.0 1.3 1.8 8.9 1.8	8.3 7.0 3 1.3	6.2 4.5 20 22 1.6
 *	RT LDG ABS	100.0 100.0 2 3 0.0 .999	52.2 87.9 33 35.7	100.0 95.7 1 4.3	100.0	78.6 93.9 8 10 15.3
	GDU SLK ABS	1348 10 1385 10 48 54 37 300#	1389 1411 17 17 22 .038+	1380 1 1432 13 16 52 502#	1426 1 1463 1 16 16 37 3007#	1368 1399 124 137 31
	GDU SHD ABS	1309 1353 49 55 44 .000#	1343 : 1384 : 20	1345 1397 13 16 52 .005#	1364 1433 16 16 69	1325 1369 128 143 44
	EST CNT ABS	26.3 26.4 38 65 0.1	39.0 43.0 26 35 4.0	29.8 31.8 10 14 2.0 .003#	41.8 42.5 13 13 0.7	32.6 33.9 116 159 1.3
	SDG VGR ABS	6.0 5.0 23 25 1.0	6.5 6.0 7 7 8.3 8.3 8.4	6.2 6.2 6.2 1.0 1.0 1.3	5.3 4.9 7 7 7 0.4	5.5 4.9 59 65 0.7
	GQU /HA %MN	103 118 3 6 15	119 100 9 18 19	121 106 4 7 15	145 123 4 22 619	122 109 20 20 35 13
	GQU /HA ABS	58.2 65.2 3 7.0 681	67.1 56.7 9 18 10.4	38.3 34.0 4 4.3 4.3	38.9 34.1 44.4 650	54.4 48.9 20 35 5.4
	TST WT ABS	-, -	52.4 51.7 9 18 0.7	58.4 55.5 3 2.9 108	54.9 51.8 1.8 3.1 1.85	54.2 52.4 16 28 1.7
	MST	16.4 15.6 3 6 0.8	26.5 26.7 11 20 0.2	15.4 15.5 4 7 7 881	21.0 21.5 4 0.5 617	21.6 21.6 24 39 0.0
	BU ACR %MN	103 118 3 6 15	119 100 100 18 19 .075*	121 106 4 7 15	145 123 4 4 22 22	122 109 20 35 13
	BU ACR ABS	92.7 103.9 3 6 11.2	107.0 90.5 9 18 16.5	61.2 54.2 4 7.0 7.0	62.0 54.5 4 7.5 650	86.7 78.0 20 35 8.7 165
	# 1	H	H	H	E	E→
	I VAR	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF	1 2 LOCS REPS DIFF
	YEAR REGION VAR	SUM	SUM	SUM	SUM	TOTAL SUM
	YEAR		94	و ت	9	TOTA

RE	PRODUCE LOCALLY, Include form number and edition date on all reproductions.	FORM APP	ROVED - OMB NO. 0581-0055
	U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in accordance 1974 (5 U. S. C. 552a) and the Paperwork F	lance with the Privacy Act of
	EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determin certificate is to be issued (7 U.S.C. 2421). In until certificate is issued (7 U.S.C. 2426).	ne if a plant variety protection
1.	NAME OF APPLICANT(S) PIONEER HI-BRED INTERNATIONAL, INC.	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME PHDG1
4	.ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
	7301 NW 62 nd AVENUE	515-270-4051	515-253-2125
	P.O.BOX 85 JOHNSTON, IA 50131-0085	7. PVPO NUMBER 97001	96
8.	Does the applicant own all rights to the variety? Mark an "X" in appropriate bl	lock. If no, please explain:	YES NO
			VEC. EL NO
9.	Is the applicant (individual or company) a U.S. national or U.S. based company	ny?	YES NO
10.	If no, give name of country Is the applicant the original owner?	If no, please answer one of the followin	<u>g:</u>
	a. If original rights to variety were owned by individual(s), is (are) the origin		
	☐ YES ☐ NO	if no, give name of country	
	b. If original rights to variety were owned by a company(ies), is(are) the ori	iginal owner(s) a U.S. based company?	
	⊠ YES □ NO	If no, give name of country	
11.	Additional explanation on ownership (if needed, use reverse for extra space):		
	PHDG1 is owned by Pioneer Hi-Bred International, Inc.		
	EASE NOTE:		
Plan	t variety protection can be afforded only to owners (not licensees) who meet one of the	e following criteria:	
1.	If the rights to the variety are owned by the original breeder, that person must be a U Which affords similar protection to nationals of the U.S. for the same genus and spec	· · · · · · · · · · · · · · · · · · ·	try, or national of a country
2.	If the rights to the variety are owned by the company which employed the original brocountry, or owned by national of a country which affords similar protection to national		ned by nationals of a UPOV member
3.	If the applicant is an owner who is not the original owner, both the original owner and	d the applicant must meet one of the above crite	eria.
The	original breeder/owner may be the individual or company who directed final breeding.	See section 41(a)(2) of the Plant Variety Prote	ection Act for definition.
in	ccording to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of info formation collection is 0581-0055. The time required to compete this information collection is estimater ata sources, gathering and maintaining the data needed, and completing and reviewing the collection of info	d to average 10 minutes per response, including the time	
pr	ne U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, colo ohibited bases apply to all programs). Persons with disabilities who require alternative means for communenter at 202-720-2600 (voice and TDD).	r, national origin, sex, religion, age, disability, political belie nication of program information (braille, large print, audiota	efs, and marital or familial status (Not all pe, etc.) should contact USDA's TARGET

To file a complaint, write Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD) USDA is an equal employment opportunity employer.